

2010 Role Delineation Study: Gerontological Nurse

National Survey Results



December 2010

© Copyright 2010 American Nurses Credentialing Center, All Rights Reserved

About this Report

This report pertaining to the practice of gerontological nurses was based on the results of American Nurses Credentialing Center's 2010 Role Delineation Study of Five Nursing Specialties: Cardiac Vascular, Gerontological, Medical-Surgical, Pediatric, and Psychiatric and Mental Health.

Prepared by:
Castle Worldwide, Inc.
900 Perimeter Park Drive, Suite G
Morrisville, NC 27560

TABLE OF CONTENTS

| | |
|--|------------|
| ACKNOWLEDGEMENTS | 4 |
| BACKGROUND | 6 |
| Role Delineation Study Overview | 6 |
| Updated Test Content Outlines | 6 |
| Role of the Content Expert Panels | 6 |
| SURVEY METHODOLOGY | 7 |
| Survey Chronology | 7 |
| Sample Selection | 7 |
| Survey Development and Measures | 7 |
| Data Collection | 8 |
| Data Analysis | 9 |
| SURVEY RESULTS | 11 |
| Demographic Information | 11 |
| Task Descriptions | 11 |
| APPENDICIES | |
| Work Activity Statements | Appendix A |
| Demographic Data Summary | Appendix B |
| Summary Statistics: Consequence, Frequency, and Performance Expectation Ratings | Appendix C |
| Task Statements Sorted by Relative Weight | Appendix D |

Acknowledgements

The American Nurses Credentialing Center (ANCC) wishes to thank a number of content experts who served on various ANCC Nursing Specialty Role Delineation Study panels for sustaining this effort and producing a role delineation study of such high caliber. Without their numerous hours of input and feedback, the study would not have been possible.

Cardiac Vascular Nurse

- Aida Ramos, MSN, RN-BC, NP-C
- Cari Zhang, MSN, RN-BC
- Mary Paquette, MSN, RN-BC
- Mary Sieggreen, MSN, RN-BC
- Rene Wells Lavoie, BSN, RN-BC
- Jimmy Callicutt, MSN, RN-BC
- Jean Vreeland, BSN, RN-BC
- Michele Sepe, MSN, ANP-BC, RN-BC
- Angela Klein, BSN, RN-BC
- Myra Torres, BSN, RN-BC
- Catherine Ware, BSN, RN-BC
- Lola Coke, PhD, ACNS-BC, RN-BC

Gerontological Nurse

- **Susan Shorter, ADN, RN-BC**
- **Leslie Green, MSN, RN-BC**
- **Camilla Camburn, MSN, RN-BC**
- **Kathy Urban, MN, RN-BC, ACNS-BC**
- **Deborah Karas, MSN, RN-BC**
- **Helen Farrar, MS, RN-BC**
- **Joseph Osentoski, BAS, RN-BC**
- **Teresa Novy, BSN, RN-BC**
- **Anastasia Reddick, BSN, RN-BC**
- **Elaine Melia, BSN, RN-BC**
- **Paula Newinski, BSN, RN-BC**
- **Darlene Shoemaker, MS, RN-BC**

Medical-Surgical Nurse

- Jane Hawks, PhD, RN-BC
- Judith Jefferson Gordon, MBA, BSN, BS, RN-BC
- Rebecca Pierce, MA, ADN, RN-BC
- Adlene McElroy, MSN, MS, RN-BC
- Vallye Brooks, BSN, RN-BC
- Donna Burns, MBA, MS, BSN, RN-BC
- Melissa Buchanan, ADN, RN-BC
- Sherlyn Farrish-Barner, MSN, RN-BC
- Kathleen Taylor, BSN, RN-BC
- Sheri Wilson, BSN, RN-BC

Pediatric Nurse

- Bonnie Brown, MSN, RN-BC
- Linda Jamison, RN-BC
- Mary Koehler, MSN, RN-BC
- Diane Saylor, RN-BC
- La Donna Northington, DNS, RN-BC
- Suzanne Wright Warner, BNS, RN-BC
- Ruth Bindler, PhD, BSN, RN-BC, DPNAP
- Geraldine Stanton, ASN, RN-BC
- Karen Leif, MA, BSN, RN-BC
- Desiree Sana-Hanscom, MHA, BSN, RN-BC

Psychiatric and Mental Health Nurse

- Duane Anderson, BSN, RN-BC
- Marlene Allen, ADN, RN-BC
- Tina Aown, MSN, RN-BC
- Susan Phillips, MSN, PMHCNS-BC, PMHNP-BC
- Catherine Meliniotis, BSN, RN-BC
- Heidi Kraft, BSN, RN-BC
- Heidi McMullan, BSN, RN-BC
- Marie Adams, BSN, RN-BC

We also would like to thank the ANCC and Castle staff who spent numerous hours working to make this study possible:

- Christine DePascale, MS, ANCC
- David Paulson, PhD, CAE, ANCC
- Gossie Nworu, MA, ANCC
- Cheray Jones, ANCC
- Diane Talley, MA, Castle (Project Manager)
- James Henderson, PhD, Castle
- James Penny, PhD, Castle

Background

The American Nurses Credentialing Center (ANCC), which was incorporated in 1991 as a subsidiary of the American Nurses Association, is the largest nursing credentialing organization in the United States. Its vision is to be a galvanizing force for quality healthcare through credentialing excellence. Currently, ANCC offers 25 examinations at various levels including diploma and associate degree, baccalaureate, and advanced practice for nurses, clinical nurse specialists, and other disciplines. Approximately 10,000 – 12,000 candidates take an ANCC certification examination each year. In addition to certification, ANCC provides services such as the Magnet and Pathways to Excellence Recognition Programs for hospitals and other facilities that demonstrate excellence in nursing services, accreditation of continuing education programs, education and consultation services, and outreach to nursing organizations around the globe.

Role Delineation Study Overview

Role delineation or job analysis studies are typically carried out at the national level with the goal of describing current practice expectations, performance requirements, and environments. ANCC has a current goal of conducting a role delineation study of nurses approximately every three years in order to capture changes in work activities (referred to as “tasks” in this report) and the knowledge and skill areas required to perform those tasks. The findings are used to update the content of its respective certification examinations. This 2010 role delineation study (performed concurrently for five nursing specialties—cardiac vascular, gerontological, medical-surgical, pediatric, and psychiatric and mental health) involved two sets of processes or activities that ran more or less concurrently: a national Web-based survey and a linking activity. The national survey was designed to collect information on the tasks that nurses working in these specialties actually perform in practice, while the linking activity identifies the major knowledge and skill areas required to perform the tasks listed in the survey. The results of both of these processes were used in the updating of the test content outlines for each examination contained within the study.

Updated Test Content Outlines

The results of this role delineation study were used to update the test content outlines for each of the five specialty examinations contained within the study. Examination forms based on this updated test content outline for gerontological nursing are scheduled to go into effect May 6, 2012. A copy of the test content outline is available on the American Nurses Credentialing Center website.

Role of the Content Expert Panels

For this study, ANCC invited content experts to develop a list of tasks and demographic items for the survey, link knowledge and skill areas to the tasks list, and finalize the test content outlines for the respective certification examinations. All of the content experts serving on the panels were certified by ANCC in the nursing specialty they represented and were invited to serve on the panels based upon expertise in their specialties.

Survey Methodology

The purpose of the development and administration of the national survey was to collect information on the tasks nurses actually perform in practice. Since the survey instrument used in this study was designed to be used across five nursing specialties, representatives from each of these five nursing specialties were asked to serve as members of a 10-member initial study workgroup (i.e., *joint role delineation panel*) that acted as a sort of steering committee for the specialty-specific role delineation panels. The members of the joint panel met for three days (February 10-12, 2010) to construct a comprehensive list of tasks to be included in the survey. The list of tasks were organized and grouped into domains. Furthermore, the joint panel also constructed a generic map of knowledge and skill areas relevant to the tasks included in the survey.

Survey Chronology

The survey development and administration timeline was as follows:

February - May 2010

- The joint role delineation panel along with staff from ANCC and Castle drafted the survey, including demographic questions and the list of tasks to be rated.
- The survey was pilot tested.

June - August 2010

- The survey was revised based on pilot study results and feedback from a few selected specialty organizations.
- The final survey was approved and administered to the national sample on the Web.

September - November 2010

- The survey task results were analyzed, and task weights were determined.
- Each specialty-specific panel met to review the survey results and task weights.

Sample Selection

In January 2010, there were a total of 2,213 active ANCC-certified gerontological nurses who had received their certification as a gerontological nurse within the previous 10 years. A random sample of 1,532 gerontological nurses was selected from the ANCC certification database, with the constraint that the participants were to have received their ANCC certification within the previous 10 years.

Survey Development and Measures

On February 10-12, 2010, the joint role delineation panel, along with members of ANCC and Castle, met in Silver Spring, Maryland, to construct the list of tasks to be included in the 2010 Role Delineation Study of five nursing specialties—cardiac vascular, gerontological, medical-surgical, pediatric, and psychiatric and mental health nursing. The panel members first reviewed and updated the target population statements for each of the five nursing specialty areas. They used the scope and standards of practice for each of these specialties, which the panelists reviewed and discussed during the meeting. From the scope and standards, the panelists identified the performance domains and tasks potentially performed by nurses working within these specialties. The goal of this process was to create a comprehensive list of relevant tasks that were potentially performed by nurses in any one of the specialty areas, regardless of whether it was performed in the others.

As a result of this meeting, the panel reached consensus on a list of 51 tasks to be used in the 2010 survey. These tasks were divided into seven domains: (1) Assessment and Diagnosis, (2) Planning and Outcomes Identification, (3) Implementation, (4) Evaluation, (5) Nurse-Patient Relationship, (6) Patient, Family, Significant Other, and Caregiver Education, and (7) Management and Leadership. The complete text of the task list is presented in **Appendix A**, including two tasks that were added to that list at a later date. The joint panel also initially identified a set of 19 demographic items for inclusion in the

survey; one was later removed based on a review of the pilot survey results. (See **Appendix B** for the final set of demographic items).

During the same meeting in February, the joint panel reviewed and approved three scales that respondents would use to rate the tasks listed in the survey — frequency (the frequency with which a task is performed), performance expectation (how soon on the job the performance of a task is expected), and consequence (the consequence of performing a task incorrectly). These three scales and their corresponding rating descriptions are illustrated in Table 1.

Table 1. Rating Scales for Rating Task Statements

| |
|--|
| <p><i>Frequency:</i> Frequency refers to how often the <specialty> nurse performs the task, considering a one-year period. The following scale was used to record frequency:</p> <ul style="list-style-type: none">0 = Never1 = Rarely2 = Sometimes3 = Often4 = Repeatedly <p><i>Performance Expectation:</i> Performance expectation refers to the point in the <specialty> nurse’s career at which he or she is expected to perform the task. The following scale was used to record performance expectation:</p> <ul style="list-style-type: none">0 = Not at all1 = Within the first six months (including exactly 6 months) of working as a <Specialty> nurse2 = After the first six months (excluding exactly 6 months) of working as a <Specialty> nurse <p><i>Consequence:</i> Consequence refers to the degree to which the inability of the <specialty> nurse to perform the task would be seen as causing harm to stakeholders. (Harm may be seen as physical, psychological, emotional, legal, financial, etc.) The following scale was used to record consequence:</p> <ul style="list-style-type: none">0 = No harm1 = Minimal harm2 = Moderate harm3 = Substantial harm4 = Extreme harm |
|--|

Members from ANCC and Castle discussed the methodology that would be used to rank the overall importance of the tasks listed in the survey. A series of analyses was used to determine the relative importance of the task statements. A more detailed description of these analyses is provided below (see the Data Analysis section), but the primary purpose was to use the consequence and frequency ratings to compute the relative importance of each task. Tasks with large relative importance scores can be interpreted as activities that are performed frequently or in which failure to perform them can result in serious harm, or both. These tasks were viewed as more critical to the certification process and were therefore weighted more heavily during the development of the updated test content outline.

In addition to computing the relative importance values for the task statements, the gerontological role delineation panel (see page 4) reviewed the performance expectation ratings and identified tasks that were either not expected of a gerontological nurse (i.e., rating of “0”) or that were expected to be performed, but not within the first six months (i.e., rating of “2”) of working in the specialty. The panel discussed each identified task to determine whether it should be removed from testing consideration and not included in the development of the updated test content outline. Final determinations for excluding a task were made based on panel consensus consistent with supporting survey data.

Data Collection

Pilot Testing. Using the same procedures for administering the national data collection, the survey was piloted in May 2010. Fifty (50) ANCC certified nurses from each specialty who had received their certification within the previous 10 years were randomly selected to take the pilot survey (250 invitations total). Overall, 84 (34 percent) of the nurses invited to take the pilot survey responded, and 22 (44 percent) of the 50 gerontological nurses responded. Also during this time, a few selected specialty organizations were asked to review the list of tasks.

Survey Revision. Two small revisions were made to the survey before it was finalized and administered to the national sample. First, after reviewing the results from the pilot survey, members from the joint panel agreed to remove one demographic question due to inconsistency of response data and low applicability to the overall purpose of the study. Second, based on feedback from the specialty organizations' reviews of the survey, two task statements were added to the survey (in Domain 1: Assessment and Diagnosis):

- Obtain a home medication list
- Perform a medication reconciliation

National Survey. On July 12, 2010, the 1,532 gerontological nurses selected to take the national Web-based survey were sent an alert letter via the United States Postal Service. On July 13, 2010, each nurse was also sent an e-mail containing the same information. The alert letter and e-mail explained the purpose and importance of the study, the eligibility criteria of the study, and stated how to access the survey via the Internet. Both indicated that the participant's responses would be kept confidential and also notified that respondents completing the survey would receive a five-hour reduction of their continuing education requirement for their recertification.

A follow-up postcard and two follow-up e-mails were sent out in the ensuing weeks. The follow-up correspondence thanked recipients if they had already submitted their completed survey and encouraged them to do so if they had not already. The follow-up postcard and first follow-up e-mails were sent to all selected nurses, and the final follow-up e-mail was sent out only to those who had not yet responded to the survey. The survey was closed on August 5, 2010.

Data Analysis

The analysis of survey data consisted of two separate processes that were performed more or less concurrently. The first process consisted of the gerontological role delineation panel's review of the performance expectation ratings and the identification of tasks that are either not expected of a gerontological nurse (i.e., rating of "0") or that are expected to be performed, but not within the first six months (i.e., rating of "2"). Discussions ensued regarding the tasks with the goal of removing those that were not seen as being expected of a gerontological nurse. Tasks designated for removal were not included in the development of the test content outline.

The second process—the analysis of the consequence and frequency ratings—was used to determine the relative importance of each of the tasks. The details of those analyses are summarized below.

Factor analysis. First, a factor analysis was performed on the consequence and frequency ratings to ensure that the data were sufficiently unidimensional for conducting an Item Response Theory (IRT) analysis. The factor analysis revealed that over 38 percent of the total variance was explained by the first unrotated factor, easily surpassing Reckase's (1979) recommendation that the first unrotated principal factor of the item responses should account for over 20% of the total variance if the items are to be rescaled using the one-parameter logistic IRT model.

Reliability analysis. The rating scales were also highly reliable. Cronbach's coefficient alpha estimates for the performance expectation, consequence, and frequency scales when applied to all the data were 0.911, 0.987, and 0.950, respectively. Cronbach's coefficient alpha, a measure of internal stability, ranges in value between 0 and 1.

IRT analysis. The factor analysis showed the survey data to be sufficiently unidimensional, which led to the subsequent IRT analysis. To put the measures of consequence and frequency on the same interval scale, the consequence and frequency ratings were rescaled using Andrich's Rating Scale model (Andrich, 1978). The Rating Scale model belongs to a family of mathematical models known collectively as Item Response Theory (IRT). The IRT analysis was implemented using *Winsteps 3.70.0.5* (Linacre, 2010). These rescaled values are referred to as *item endorsements*.

Computation of Relative Importance. The consequence and frequency item endorsements that resulted from the IRT analysis were then used to determine the relative importance of each task. The consequence and frequency item endorsements were first transformed to the N(5,1) scale to eliminate negative values. Subsequently, the transformed consequence and frequency item endorsements were multiplied together, and the magnitude of those values were treated as indicators of the relative importance of each task. The relative importance of the tasks, including tasks that were eventually removed from testing consideration, is reported in **Appendix D**.

Survey Results

The total sample size of the national survey included 1,532 ANCC certified gerontological nurses who had received their certification within the previous 10 years. A total of 622 valid gerontological nurse surveys were returned for an overall response rate of 41 percent.

Demographic Information

Appendix B details the gerontological nurses' responses to the final survey's 18 demographic questions that included inquiry on the nurses' backgrounds and practice settings.

Demographic Background

Approximately 94 percent of the respondents were female and 82 percent reported to be white. Approximately 70 percent of the sample fell into the age group of 40-59 years of age.

Approximately 40 percent of the gerontological respondents indicated that a Bachelor's in Nursing was their highest degree in nursing, with another 31 and 19 percent indicating Associate in Nursing and Master's in Nursing as their highest degree, respectively. Less than one percent indicated they held a Doctorate in Nursing (Practice or Research).

The average number of years of experience the gerontological nurse respondents had as an RN was just over 21 years. The respondents also reported on average 11 years of experience working as a gerontological nurse.

Practice Settings

Approximately 38 percent of the gerontological nurse respondents indicated that they practiced in cities with populations between 50,000 and 249,999. Towns with a population between 2,500 – 49,999 had the second highest percent of respondents (28 percent).

In terms of practice setting, the highest percentage of gerontological nurse respondents indicating they practice in a long term care facility (28 percent). Approximately 16 percent listed a medical-surgical facility as their primary practice setting.

Approximately 35 percent of gerontological nurses reported that "staff nurse" best described their work. Another 17 percent listed "management" as best describing their work. Sixty-two (62) and 82 percent of respondents described their patients as dealing with acute and chronic illnesses, respectively. The gerontological nurses also indicated that almost 99 percent of their time was spent in caring for adults (18 to 65 years—24.8%) or aging adults (ages over 65 years—73.8%). When asked how many hours per week on average they spent working as a gerontological nurse, the average response was 33.6 hours with a standard deviation of 12.2 hours.

Task Descriptions

Task Summary Statistics. Descriptive statistics (means, standard deviations, and medians) for the consequence and frequency ratings and frequency statistics for the performance expectation ratings are reported in **Appendix C** for all 53 tasks.

The mode of the performance expectation ratings, the mean for the frequency and consequence ratings, and relative importance (based on the multiplied item endorsements from the IRT analysis) is reported for all 53 tasks in **Appendix D**, presented in rank order or relative importance. It should be noted that **Appendices C** and **D** contain statistics for the entire set of tasks before the gerontological panel reached a decision to remove a few of the tasks from consideration due to a variety of reasons (e.g., performance expectation ratings that were either too high or too low).

Performance Expectation. Only one task—1.12: Identify nursing diagnoses using a different system—had a performance expectation rating mode of zero (i.e., not at all). Three tasks had a rating

mode of 2 (i.e., performed after 6 months of working as a gerontological nurse)—7.4: Serve as a clinical content expert for the design and enhancement of policies, procedures, processes, and systems that affect nursing care, 7.1: Serve as a preceptor, and 6.4: Develop educational programs for groups.

Consequence. The highest rated task with respect to consequence was 3.5: Administer medications as prescribed using evidence-based, developmentally appropriate, age-appropriate techniques. The second and third rated tasks, both from the Assessment and Diagnosis Domain (Domain 1), were 1.3: Perform a medication reconciliation and 1.2: Obtain a home medication list, respectively. The lowest rated task was again 1.12: Identify nursing diagnoses using a different system.

Frequency. Three of the top four rated tasks with respect to frequency came from the Nurse-Patient Relationship domain (domain 5), and included 5.4: Maintain a developmentally appropriate therapeutic relationship, 5.2: Establish trust, and 5.3: Develop rapport. Task 7.7: Follow legal and regulatory requirements in nursing care delivery and management was the second rated task. The lowest rated task was again found to be task 1.12.

Task Relative Importance. The task with the highest relative importance value was 1.2: Obtain a home medication list. Task 1.5: Complete psychosocial assessment using age-appropriate, system-specific, evidence-based tools and assessment techniques was the second highest rated task. Not surprisingly, task 1.12: Identify nursing diagnoses using a different system was the lowest rated task with respect to relative importance.

Appendix A

Task Statements

Domain 1: Assessment and Diagnosis

Tasks:

- 1.1 Obtain patient history using age-appropriate, system-specific, evidence-based tools.
- 1.2 Obtain a home medication list.
- 1.3 Perform a medication reconciliation
- 1.4 Perform physical examination using age-appropriate, system-specific, evidence-based assessment techniques.
- 1.5 Complete psychosocial assessment using age-appropriate, system-specific, evidence-based tools and assessment techniques.
- 1.6 Collect data on psychoses.
- 1.7 Obtain diagnostic test results.
- 1.8 Review findings provided by others.
- 1.9 Synthesize available data and knowledge to identify patterns and variances.
- 1.10 Identify nursing diagnoses using the North American Nursing Diagnosis Association-International (NANDA-I) taxonomy.
- 1.11 Identify nursing diagnoses using clinical pathways.
- 1.12 Identify nursing diagnoses using a different system. [Specify the system]
- 1.13 Create a problem list based on assessment data.
- 1.14 Document assessment findings in patient records.

Domain 2: Planning and Outcomes Identification

Tasks:

- 2.1 Prioritize nursing diagnoses and/or problems.
- 2.2 Formulate expected outcomes with patient, family, significant other, and interdisciplinary team involvement using Nursing Outcomes Classification (NOC).
- 2.3 Formulate expected outcomes with patient, family, significant other, and interdisciplinary team involvement using clinical pathways or a different system. [Specify other]
- 2.4 Develop an individualized, developmentally appropriate plan of care.
- 2.5 Document plan of care and expected outcomes in patient records.

Domain 3: Implementation

Tasks:

- 3.1 Use Nursing Interventions Classification (NIC) as the basis for interventions specific to the plan of care.
- 3.2 Use other evidence-based practice guidelines as the basis for interventions specific to the plan of care.
- 3.3 Create a safe, developmentally appropriate, therapeutic environment conducive to care.
- 3.4 Coordinate patient care.
- 3.5 Administer medications as prescribed using evidence-based, developmentally appropriate, age-appropriate techniques.
- 3.6 Document nursing interventions in patient records.

Domain 4: Evaluation

Tasks:

- 4.1 Collect data related to the patient's response to interventions.
- 4.2 Collect data on factors that impact the patient's care.
- 4.3 Evaluate the patient's response to interventions and the effectiveness of the plan of care.
- 4.4 Update the plan of care.
- 4.5 Communicate changes to the patient, family, significant other, and interdisciplinary team.
- 4.6 Document the patient's response to interventions and changes to the plan of care in patient records.

Appendix A – Task Statements

2010 Role Delineation Study: Gerontological Nurse – National Survey Results

Domain 5: Nurse-Patient Relationship

Tasks:

- 5.1 Approach the patient in a developmentally appropriate manner.
- 5.2 Establish trust.
- 5.3 Develop rapport.
- 5.4 Maintain a developmentally appropriate therapeutic relationship.
- 5.5 Support the patient's parents, family, significant others, and caregivers.
- 5.6 Maintain appropriate physical and emotional boundaries.
- 5.7 Serve as patient advocate.
- 5.8 Document pertinent aspects of the nurse-patient relationship in patient records.

Domain 6: Patient, Family, Significant Other, and Caregiver Education

Tasks:

- 6.1 Identify learning needs.
- 6.2 Identify barriers to learning.
- 6.3 Develop an individualized education plan with the involvement of the patient, family, significant other, and caregiver.
- 6.4 Develop educational programs for groups.
- 6.5 Implement the education plan.
- 6.6 Evaluate the education plan's effectiveness.
- 6.7 Document the education provided and its effectiveness in patient records.

Domain 7: Management and Leadership

Tasks:

- 7.1 Serve as a preceptor.
- 7.2 Coordinate patient safety measures.
- 7.3 Improve quality of nursing care delivery.
- 7.4 Serve as a clinical content expert for the design and enhancement of policies, procedures, processes, and systems that affect nursing care.
- 7.5 Use electronic information systems in nursing care delivery.
- 7.6 Follow ethical standards in nursing care delivery and management.
- 7.7 Follow legal and regulatory requirements in nursing care delivery and management.

Appendix B

Demographic Data Summary

1: What is your gender?

| | Count | Percent |
|--------|-------|---------|
| Female | 582 | 94.0 |
| Male | 37 | 6.0 |
| Total | 619 | 100.0 |

2: What is your age?

| | Count | Percent |
|-----------------------|-------|---------|
| Under 20 years old | 0 | 0.0 |
| 20 to 29 years old | 15 | 2.4 |
| 30 to 39 years old | 79 | 12.8 |
| 40 to 49 years old | 161 | 26.1 |
| 50 to 59 years old | 268 | 43.4 |
| 60 years old and over | 94 | 15.2 |
| Total | 617 | 100.0 |

3: What is your ethnicity?

| | Count | Percent |
|---------------------------------|-------|---------|
| African-American | 35 | 5.7 |
| White, non-Hispanic | 504 | 82.0 |
| Asian/Pacific Islander | 45 | 7.3 |
| American Indian/ Alaskan Native | 2 | 0.3 |
| Hispanic/Latino | 15 | 2.4 |
| Other | 14 | 2.3 |
| Total | 615 | 100.0 |

4: Which of the following describes your entry-level (basic) education in nursing?

| | Count | Percent |
|-----------------------------|-------|---------|
| Diploma in Nursing | 108 | 17.6 |
| Associate Degree in Nursing | 256 | 41.6 |
| Bachelor's in Nursing | 221 | 35.9 |
| Other | 30 | 4.9 |
| Total | 615 | 100.0 |

5: What is the highest degree you have earned in nursing?

| | Count | Percent |
|---|-------|---------|
| Diploma in Nursing | 47 | 7.6 |
| Associate Degree in Nursing | 189 | 30.6 |
| Bachelor's in Nursing | 245 | 39.6 |
| Master's in Nursing | 116 | 18.8 |
| Doctorate in Nursing Practice (DNP) | 3 | 0.5 |
| Doctorate in Nursing Research (e.g., Ph.D., DNS, DSN) | 0 | 0.0 |
| Other | 18 | 2.9 |
| Total | 618 | 100.0 |

6: Do you hold any degrees outside of nursing?

| | Count | Percent |
|-------|-------|---------|
| Yes | 154 | 25.1 |
| No | 460 | 74.9 |
| Total | 614 | 100.0 |

7: Are you certified as a gerontological nurse?

| | Count | Percent |
|-------|-------|---------|
| Yes | 608 | 99.7 |
| No | 2 | 0.3 |
| Total | 610 | 100.0 |

8a: How many years have you been certified as a gerontological nurse?

| | Count | Percent |
|---------|-------|---------|
| 0-5 | 463 | 76.4 |
| 6-10 | 124 | 20.5 |
| 11-15 | 14 | 2.3 |
| Over 15 | 5 | 0.8 |
| Total | 606 | 100.0 |

8b: How many years have you been certified as a gerontological nurse?

| | |
|------|-------|
| N | 609 |
| Min | 0 |
| Max | 44 |
| Mean | 4.18 |
| SD | 4.035 |

9: Do you hold other nursing certifications?

| | Count | Percent |
|-------|-------|---------|
| Yes | 223 | 36.3 |
| No | 391 | 63.7 |
| Total | 614 | 100.0 |

10a: How many years have you been in practice as an RN?

| | Count | Percent |
|---------|-------|---------|
| 0-5 | 42 | 6.9 |
| 6-10 | 84 | 13.8 |
| 11-15 | 100 | 16.4 |
| 16-20 | 93 | 15.3 |
| 21-25 | 69 | 11.3 |
| 26-30 | 70 | 11.5 |
| 31-35 | 85 | 14.0 |
| 36-40 | 45 | 7.4 |
| Over 40 | 20 | 3.3 |
| Total | 608 | 100.0 |

10b: How many years have you been in practice as an RN?

| | |
|------|-------|
| N | 608 |
| Min | 3 |
| Max | 50 |
| Mean | 21.23 |
| SD | 11.13 |

11a: How many years have you worked as a gerontological nurse?

| Years | Count | Percent |
|---------|-------|---------|
| 0-5 | 225 | 36.7 |
| 6-10 | 130 | 21.2 |
| 11-15 | 97 | 15.8 |
| 16-20 | 90 | 14.7 |
| 21-25 | 31 | 5.1 |
| 26-30 | 16 | 2.6 |
| 31-35 | 18 | 2.9 |
| 36-40 | 4 | 0.7 |
| Over 40 | 2 | 0.3 |
| Total | 613 | 100.0 |

11b: How many years have you worked as a gerontological nurse?

| | |
|------|-------|
| N | 613 |
| Min | 0 |
| Max | 46 |
| Mean | 10.94 |
| SD | 8.66 |

12: How many hours per week do you work as a gerontological nurse?

| Hours | Count | Percent |
|---------|-------|---------|
| 0-20 | 101 | 16.5 |
| 21-35 | 108 | 17.7 |
| 36-40 | 339 | 55.5 |
| Over 40 | 63 | 10.3 |
| Total | 611 | 100.0 |

13: What percent of time do you spend with each type of patient?

| | Infant (pre-birth to 2 yrs) | | Child (ages 3 to 12) | | Adolescents (ages 13 to 17) | | Adults (ages 18 to 65) | | Aging Adult (over 65 yrs) | |
|-------------------|-----------------------------|---------|----------------------|---------|-----------------------------|---------|------------------------|---------|---------------------------|---------|
| | Count | Percent | Count | Percent | Count | Percent | Count | Percent | Count | Percent |
| 0 - 20% | 621 | 99.8 | 617 | 99.2 | 619 | 99.5 | 309 | 49.7 | 20 | 3.2 |
| 21 - 40% | 0 | 0 | 4 | 0.6 | 3 | 0.5 | 183 | 29.4 | 31 | 5 |
| 41 - 60% | 0 | 0 | 0 | 0 | 0 | 0 | 108 | 17.4 | 168 | 27 |
| 61 - 80% | 0 | 0 | 1 | 0.2 | 0 | 0 | 15 | 2.4 | 165 | 26.5 |
| 81 - 100% | 1 | 0.2 | 0 | 0 | 0 | 0 | 7 | 1.1 | 238 | 38.3 |
| Total | 622 | 100 | 622 | 100 | 622 | 100 | 622 | 100 | 622 | 100 |
| Average (percent) | 0.3 | | 0.5 | | 0.6 | | 24.8 | | 73.8 | |

14: Which of the following best describes your work?

| | Count | Percent |
|--|-------|---------|
| Staff Nurse (including community, clinic, in-patient/client, or other setting) | 217 | 35.3 |
| Clinical Nurse Specialist | 11 | 1.8 |
| Nurse Practitioner | 14 | 2.3 |
| Case Manager | 56 | 9.1 |
| Education | 44 | 7.2 |
| Management | 102 | 16.6 |
| Research | 3 | 0.5 |
| Charge Nurse | 81 | 13.2 |
| Clinical Nurse | 4 | 0.7 |
| Other | 82 | 13.4 |
| Total | 614 | 100.0 |

15: Which of the following best describes your primary practice setting?

| | Count | Percent |
|--|-------|---------|
| Medical Unit | 65 | 10.6 |
| Surgical Unit | 12 | 1.9 |
| Medical/Surgical | 98 | 15.9 |
| Intensive Care Unit | 11 | 1.8 |
| Cardiac Surgery Intensive Care Unit | 1 | 0.2 |
| Correctional care Unit | 0 | 0.0 |
| Same Day Surgery | 2 | 0.3 |
| Same Day Medical | 1 | 0.2 |
| Recovery Room | 5 | 0.8 |
| Labor and Delivery/Post Partum | 1 | 0.2 |
| Emergency Department | 3 | 0.5 |
| Outpatient Clinic | 21 | 3.4 |
| Home care or Home Health | 63 | 10.2 |
| Hospice | 7 | 1.1 |
| Long Term Care | 165 | 26.8 |
| Psychiatric/Mental Health Facility | 17 | 2.8 |
| Pediatrics | 0 | 0.0 |
| Rehabilitation | 34 | 5.5 |
| School (Elementary, Middle, or High) | 1 | 0.2 |
| College or Post-Secondary Education Facility | 10 | 1.6 |
| Pediatric Intensive Care Unit | 0 | 0.0 |
| Neonatal Intensive Care Unit | 0 | 0.0 |
| Other | 99 | 16.1 |
| Total | 616 | 100.0 |

16: Which of the following describes your patients (more than one could be selected)

| | n | number selecting | percent |
|----------------------|-----|------------------|---------|
| Well (minor illness) | 622 | 78 | 12.5 |
| Maternity | 622 | 10 | 1.6 |
| Acute | 622 | 384 | 61.7 |
| Chronic | 622 | 507 | 81.5 |
| Terminally Ill | 622 | 282 | 45.3 |
| Psychiatric | 622 | 281 | 45.2 |
| Other | 622 | 45 | 7.2 |

17: In which state is your primary practice setting located?

| | Count | Percent |
|---------------------------|-------|---------|
| Alabama (AL) | 4 | 0.6 |
| Alaska (AK) | 4 | 0.6 |
| Arizona (AZ) | 7 | 1.1 |
| Arkansas (AR) | 3 | 0.5 |
| California (CA) | 28 | 4.5 |
| Colorado (CO) | 2 | 0.3 |
| Connecticut (CT) | 15 | 2.4 |
| Delaware | 19 | 3.1 |
| District of Columbia (DC) | 1 | 0.2 |
| Florida (FL) | 29 | 4.7 |
| Georgia (GA) | 9 | 1.4 |
| Guam (GU) | 0 | 0.0 |
| Hawaii | 3 | 0.5 |
| Idaho | 1 | 0.2 |
| Illinois (IL) | 20 | 3.2 |
| Indiana (IN) | 4 | 0.6 |
| Iowa (IA) | 13 | 2.1 |
| Kansas (KS) | 4 | 0.6 |
| Kentucky (KY) | 4 | 0.6 |
| Louisiana (LA) | 7 | 1.1 |
| Maine (ME) | 7 | 1.1 |
| Maryland (MD) | 3 | 0.5 |
| Massachusetts (MA) | 27 | 4.3 |
| Michigan (MI) | 22 | 3.5 |
| Minnesota (MN) | 14 | 2.3 |
| Mississippi (MS) | 4 | 0.6 |
| Missouri (MO) | 6 | 1.0 |
| Montana (MT) | 0 | 0.0 |
| Nebraska (NE) | 24 | 3.9 |
| Nevada (NV) | 0 | 0.0 |
| New Hampshire (NH) | 7 | 1.1 |
| New Jersey (NJ) | 47 | 7.6 |
| New Mexico (NM) | 2 | 0.3 |
| New York (NY) | 62 | 10.0 |
| North Carolina (NC) | 14 | 2.3 |
| North Dakota (ND) | 6 | 1.0 |
| Ohio (OH) | 21 | 3.4 |
| Oklahoma (OK) | 6 | 1.0 |
| Oregon (OR) | 11 | 1.8 |
| Pennsylvania (PA) | 40 | 6.4 |
| Rhode Island (RI) | 3 | 0.5 |
| South Carolina (SC) | 8 | 1.3 |
| South Dakota (SD) | 12 | 1.9 |
| Tennessee (TN) | 14 | 2.3 |
| Texas (TX) | 12 | 1.9 |
| Utah (UT) | 0 | 0.0 |
| Vermont (VT) | 1 | 0.2 |

17: In which state is your primary practice setting located?

| | Count | Percent |
|--------------------|------------|--------------|
| Virginia (VA) | 12 | 1.9 |
| Washington (WA) | 4 | 0.6 |
| West Virginia (WV) | 2 | 0.3 |
| Wisconsin (WI) | 5 | 0.8 |
| Wyoming (WY) | 2 | 0.3 |
| Missing | 47 | 7.6 |
| Total | 622 | 100.0 |

18: What is the geographical location of your practice setting?

| | Count | Percent |
|---|------------|--------------|
| Rural (population less than 2,500) | 29 | 4.7 |
| Town (population 2,500 to 49,999) | 167 | 27.3 |
| City (population 50,000 to 249,999) | 231 | 37.7 |
| Metropolitan (population 25,000 to 999,999) | 101 | 16.5 |
| Greater Metropolitan (population greater than 999,999) | 63 | 10.3 |
| Regionally (population across a designated area such as several states) | 18 | 2.9 |
| Nationally (population across the United States) | 2 | 0.3 |
| Internationally (population across multiple nations) | 1 | 0.2 |
| Total | 612 | 100.0 |

Appendix C

Summary Statistics: Consequence, Frequency, and Performance Expectation Ratings

Note: For each of the tables in Appendix C, recall the following descriptions for each of the three rating scales:

Consequence: Consequence refers to the degree to which the inability of the <specialty> nurse to perform duties in each performance domain or task would be seen as causing harm to stakeholders. (Harm may be seen as physical, psychological, emotional, legal, financial, etc.) The following scale was used to record consequence:

- 0 = No harm
- 1 = Minimal harm
- 2 = Moderate harm
- 3 = Substantial harm
- 4 = Extreme harm

Frequency: Frequency refers to how often the <specialty> nurse performs duties in each of the performance domains or tasks considering a one-year period. The following scale was used to record frequency:

- 0 = Never
- 1 = Rarely
- 2 = Sometimes
- 3 = Often
- 4 = Repeatedly

Performance Expectation: Performance expectation refers to the point in the certified nurse's career at which he or she is expected to perform the work activity. The following scale was used to record performance expectation:

- 0 = Not at all
- 1 = Within the first six months (including exactly 6 months) of working as a <Specialty> nurse
- 2 = After the first six months (excluding exactly 6 months) of working as a <Specialty> nurse

Domain 1 Ratings: Assessment and Diagnosis

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|---|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| 1.1: Obtain patient history using age-appropriate, system-specific, evidence-based tools. | 619 | 2.0 | 1.8 | 1.2 | 619 | 4.0 | 3.3 | 0.9 | 0 | 11 | 1.8 |
| | | | | | | | | | 1 | 601 | 96.9 |
| | | | | | | | | | 2 | 8 | 1.3 |
| 1.2: Obtain a home medication list. | 618 | 3.0 | 2.5 | 1.3 | 619 | 4.0 | 3.3 | 0.9 | 0 | 15 | 2.4 |
| | | | | | | | | | 1 | 598 | 96.5 |
| | | | | | | | | | 2 | 7 | 1.1 |
| 1.3: Perform a medication reconciliation | 619 | 3.0 | 2.5 | 1.3 | 619 | 4.0 | 3.3 | 1.0 | 0 | 21 | 3.4 |
| | | | | | | | | | 1 | 575 | 92.7 |
| | | | | | | | | | 2 | 24 | 3.9 |
| 1.4: Perform physical examination using age-appropriate, system-specific, evidence-based assessment techniques. | 618 | 2.0 | 2.1 | 1.2 | 619 | 4.0 | 3.3 | 0.9 | 0 | 16 | 2.6 |
| | | | | | | | | | 1 | 558 | 90.1 |
| | | | | | | | | | 2 | 45 | 7.3 |
| 1.5: Complete psychosocial assessment using age-appropriate, system-specific, evidence-based tools and assessment techniques. | 619 | 2.0 | 1.7 | 1.1 | 619 | 3.0 | 3.1 | 1.0 | 0 | 19 | 3.1 |
| | | | | | | | | | 1 | 534 | 86.3 |
| | | | | | | | | | 2 | 66 | 10.7 |
| 1.6: Collect data on psychoses. | 619 | 2.0 | 1.7 | 1.2 | 619 | 2.0 | 2.3 | 1.2 | 0 | 79 | 12.8 |
| | | | | | | | | | 1 | 398 | 64.3 |
| | | | | | | | | | 2 | 142 | 22.9 |
| 1.7: Obtain diagnostic test results. | 618 | 2.0 | 2.2 | 1.3 | 617 | 3.0 | 3.2 | 0.9 | 0 | 14 | 2.3 |
| | | | | | | | | | 1 | 577 | 93.4 |
| | | | | | | | | | 2 | 27 | 4.4 |
| 1.8: Review findings provided by others. | 618 | 2.0 | 1.8 | 1.2 | 618 | 3.0 | 3.1 | 0.9 | 0 | 13 | 2.1 |
| | | | | | | | | | 1 | 549 | 88.8 |
| | | | | | | | | | 2 | 56 | 9.1 |
| 1.9: Synthesize available data and knowledge to identify patterns and variances. | 618 | 2.0 | 1.7 | 1.1 | 619 | 3.0 | 2.7 | 1.1 | 0 | 36 | 5.8 |
| | | | | | | | | | 1 | 356 | 57.5 |
| | | | | | | | | | 2 | 227 | 36.7 |
| 1.10: Identify nursing diagnoses using the North American Nursing Diagnosis Association- | 618 | 1.0 | 1.2 | 1.1 | 619 | 3.0 | 2.6 | 1.3 | 0 | 77 | 12.4 |
| | | | | | | | | | 1 | 476 | 76.9 |

* Consequence Ratings: 0 = No Harm, 1 = Minimal Harm, 2 = Moderate Harm, 3 = Substantial Harm, 4 = Extreme Harm. Frequency Ratings: 0 = Never, 1 = Rarely, 2 = Sometimes, 3 = Often, 4 = Repeatedly. Performance Expectation Ratings: 0 = Not at all; 1 = Within the first 6 months of working as a <specialty> nurse; 2 = After the first 6 months of working as a <specialty> nurse.

Appendix C – Summary Statistics: Consequence, Frequency, and Performance Expectation Ratings 2010 Role Delineation Study: Gerontological Nurse – National Survey Results

© Copyright 2010 American Nurses Credentialing Center, All Rights Reserved

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|---|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| International (NANDA-I) taxonomy. | | | | | | | | | 2 | 66 | 10.7 |
| 1.11: Identify nursing diagnoses using clinical pathways. | 618 | 1.0 | 1.3 | 1.1 | 619 | 3.0 | 2.6 | 1.2 | 0 | 67 | 10.8 |
| | | | | | | | | | 1 | 480 | 77.5 |
| | | | | | | | | | 2 | 72 | 11.6 |
| 1.12: Identify nursing diagnoses using a different system. [Specify the system] | 615 | 0.0 | 0.7 | 1.0 | 616 | 1.0 | 1.3 | 1.5 | 0 | 326 | 52.8 |
| | | | | | | | | | 1 | 240 | 38.9 |
| | | | | | | | | | 2 | 51 | 8.3 |
| 1.13: Create a problem list based on assessment data. | 619 | 2.0 | 1.6 | 1.1 | 620 | 3.0 | 3.0 | 1.1 | 0 | 33 | 5.3 |
| | | | | | | | | | 1 | 541 | 87.4 |
| | | | | | | | | | 2 | 45 | 7.3 |
| 1.14: Document assessment findings in patient records. | 619 | 2.0 | 2.1 | 1.3 | 619 | 4.0 | 3.6 | 0.7 | 0 | 9 | 1.5 |
| | | | | | | | | | 1 | 601 | 97.1 |
| | | | | | | | | | 2 | 9 | 1.5 |

Domain 2 Ratings: Planning and Outcomes Identification

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|---|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| 2.1: Prioritize nursing diagnoses and/or problems. | 619 | 2.0 | 1.8 | 1.1 | 619 | 4.0 | 3.3 | 0.9 | 0 | 14 | 2.3 |
| | | | | | | | | | 1 | 551 | 89.0 |
| | | | | | | | | | 2 | 54 | 8.7 |
| 2.2: Formulate expected outcomes with patient, family, significant other, and interdisciplinary team involvement using Nursing Outcomes Classification (NOC). | 619 | 1.0 | 1.4 | 1.1 | 619 | 3.0 | 2.7 | 1.2 | 0 | 72 | 11.6 |
| | | | | | | | | | 1 | 430 | 69.5 |
| | | | | | | | | | 2 | 117 | 18.9 |
| 2.3: Formulate expected outcomes with patient, family, significant other, and interdisciplinary team involvement using clinical pathways or a different system. | 617 | 1.0 | 1.3 | 1.1 | 617 | 3.0 | 2.6 | 1.3 | 0 | 84 | 13.6 |
| | | | | | | | | | 1 | 421 | 68.2 |
| | | | | | | | | | 2 | 112 | 18.2 |
| 2.4: Develop an individualized, developmentally appropriate plan of care. | 618 | 2.0 | 1.7 | 1.1 | 618 | 3.0 | 3.1 | 0.9 | 0 | 8 | 1.3 |
| | | | | | | | | | 1 | 541 | 87.5 |

* Consequence Ratings: 0 = No Harm, 1 = Minimal Harm, 2 = Moderate Harm, 3 = Substantial Harm, 4 = Extreme Harm. Frequency Ratings: 0 = Never, 1 = Rarely, 2 = Sometimes, 3 = Often, 4 = Repeatedly. Performance Expectation Ratings: 0 = Not at all; 1 = Within the first 6 months of working as a <specialty> nurse; 2 = After the first 6 months of working as a <specialty> nurse.

Appendix C – Summary Statistics: Consequence, Frequency, and Performance Expectation Ratings 2010 Role Delineation Study: Gerontological Nurse – National Survey Results

© Copyright 2010 American Nurses Credentialing Center, All Rights Reserved

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|--|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| | | | | | | | | | 2 | 69 | 11.2 |
| 2.5: Document plan of care and expected outcomes in patient records. | 617 | 2.0 | 1.7 | 1.1 | 618 | 3.0 | 3.3 | 0.9 | 0 | 13 | 2.1 |
| | | | | | | | | | 1 | 568 | 91.9 |
| | | | | | | | | | 2 | 37 | 6.0 |

Domain 3 Ratings: Implementation

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|--|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| 3.1: Use Nursing Interventions Classification (NIC) as the basis for interventions specific to the plan of care. | 616 | 1.0 | 1.3 | 1.1 | 615 | 3.0 | 2.4 | 1.5 | 0 | 122 | 19.8 |
| | | | | | | | | | 1 | 418 | 67.9 |
| | | | | | | | | | 2 | 76 | 12.3 |
| 3.2: Use other evidence-based practice guidelines as the basis for interventions specific to the plan of care. | 615 | 2.0 | 1.5 | 1.1 | 615 | 3.0 | 2.9 | 1.1 | 0 | 36 | 5.9 |
| | | | | | | | | | 1 | 478 | 77.7 |
| | | | | | | | | | 2 | 101 | 16.4 |
| 3.3: Create a safe, developmentally appropriate, therapeutic environment conducive to care. | 616 | 2.0 | 2.2 | 1.3 | 616 | 4.0 | 3.5 | 0.8 | 0 | 4 | 0.6 |
| | | | | | | | | | 1 | 583 | 94.6 |
| | | | | | | | | | 2 | 29 | 4.7 |
| 3.4: Coordinate patient care. | 616 | 2.0 | 2.0 | 1.2 | 616 | 4.0 | 3.5 | 0.8 | 0 | 8 | 1.3 |
| | | | | | | | | | 1 | 527 | 85.6 |
| | | | | | | | | | 2 | 81 | 13.1 |
| 3.5: Administer medications as prescribed using evidence-based, developmentally appropriate, age-appropriate techniques. | 616 | 3.0 | 2.9 | 1.4 | 615 | 4.0 | 3.6 | 0.9 | 0 | 22 | 3.6 |
| | | | | | | | | | 1 | 583 | 94.6 |
| | | | | | | | | | 2 | 11 | 1.8 |
| 3.6: Document nursing interventions in patient records. | 615 | 2.0 | 1.9 | 1.3 | 616 | 4.0 | 3.6 | 0.8 | 0 | 14 | 2.3 |
| | | | | | | | | | 1 | 597 | 96.9 |
| | | | | | | | | | 2 | 5 | 0.8 |

* Consequence Ratings: 0 = No Harm, 1 = Minimal Harm, 2 = Moderate Harm, 3 = Substantial Harm, 4 = Extreme Harm. Frequency Ratings: 0 = Never, 1 = Rarely, 2 = Sometimes, 3 = Often, 4 = Repeatedly. Performance Expectation Ratings: 0 = Not at all; 1 = Within the first 6 months of working as a <specialty> nurse; 2 = After the first 6 months of working as a <specialty> nurse.

Appendix C – Summary Statistics: Consequence, Frequency, and Performance Expectation Ratings 2010 Role Delineation Study: Gerontological Nurse – National Survey Results

Domain 4 Ratings: Evaluation

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|---|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| 4.1: Collect data related to the patient's response to interventions. | 616 | 2.0 | 1.8 | 1.2 | 615 | 4.0 | 3.3 | 1.0 | 0 | 16 | 2.6 |
| | | | | | | | | | 1 | 549 | 89.1 |
| | | | | | | | | | 2 | 51 | 8.3 |
| 4.2: Collect data on factors that impact the patient's care. | 615 | 2.0 | 1.8 | 1.2 | 615 | 3.0 | 3.1 | 1.0 | 0 | 24 | 3.9 |
| | | | | | | | | | 1 | 525 | 85.4 |
| | | | | | | | | | 2 | 66 | 10.7 |
| 4.3: Evaluate the patient's response to interventions and the effectiveness of the plan of care. | 615 | 2.0 | 1.9 | 1.2 | 615 | 4.0 | 3.4 | 0.8 | 0 | 10 | 1.6 |
| | | | | | | | | | 1 | 567 | 92.2 |
| | | | | | | | | | 2 | 38 | 6.2 |
| 4.4: Update the plan of care. | 615 | 2.0 | 1.7 | 1.1 | 615 | 3.0 | 3.3 | 0.8 | 0 | 8 | 1.3 |
| | | | | | | | | | 1 | 568 | 92.4 |
| | | | | | | | | | 2 | 39 | 6.3 |
| 4.5: Communicate changes to the patient, family, significant other, and interdisciplinary team. | 615 | 2.0 | 2.0 | 1.2 | 615 | 4.0 | 3.5 | 0.7 | 0 | 1 | 0.2 |
| | | | | | | | | | 1 | 591 | 96.1 |
| | | | | | | | | | 2 | 23 | 3.7 |
| 4.6: Document the patient's response to interventions and changes to the plan of care in patient records. | 615 | 2.0 | 1.9 | 1.2 | 614 | 4.0 | 3.5 | 0.8 | 0 | 5 | 0.8 |
| | | | | | | | | | 1 | 585 | 95.1 |
| | | | | | | | | | 2 | 25 | 4.1 |

Domain 5 Ratings: Nurse-Patient Relationship

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|--|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| 5.1: Approach the patient in a developmentally appropriate manner. | 614 | 2.0 | 1.6 | 1.1 | 614 | 4.0 | 3.7 | 0.6 | 0 | 1 | 0.2 |
| | | | | | | | | | 1 | 595 | 96.9 |
| | | | | | | | | | 2 | 18 | 2.9 |
| 5.2: Establish trust. | 614 | 2.0 | 1.8 | 1.2 | 614 | 4.0 | 3.7 | 0.6 | 0 | 0 | 0.0 |
| | | | | | | | | | 1 | 604 | 98.4 |
| | | | | | | | | | 2 | 10 | 1.6 |

* Consequence Ratings: 0 = No Harm, 1 = Minimal Harm, 2 = Moderate Harm, 3 = Substantial Harm, 4 = Extreme Harm. Frequency Ratings: 0 = Never, 1 = Rarely, 2 = Sometimes, 3 = Often, 4 = Repeatedly. Performance Expectation Ratings: 0 = Not at all; 1 = Within the first 6 months of working as a <specialty> nurse; 2 = After the first 6 months of working as a <specialty> nurse.

Appendix C – Summary Statistics: Consequence, Frequency, and Performance Expectation Ratings 2010 Role Delineation Study: Gerontological Nurse – National Survey Results

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|---|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| 5.3: Develop rapport. | 612 | 2.0 | 1.6 | 1.2 | 612 | 4.0 | 3.7 | 0.6 | 0 | 2 | 0.3 |
| | | | | | | | | | 1 | 596 | 97.2 |
| | | | | | | | | | 2 | 15 | 2.4 |
| 5.4: Maintain a developmentally appropriate therapeutic relationship. | 614 | 2.0 | 2.0 | 1.3 | 614 | 4.0 | 3.7 | 0.6 | 0 | 2 | 0.3 |
| | | | | | | | | | 1 | 599 | 97.6 |
| | | | | | | | | | 2 | 13 | 2.1 |
| 5.5: Support the patient's parents, family, significant others, and caregivers. | 614 | 2.0 | 1.6 | 1.1 | 614 | 4.0 | 3.6 | 0.7 | 0 | 6 | 1.0 |
| | | | | | | | | | 1 | 572 | 93.2 |
| | | | | | | | | | 2 | 36 | 5.9 |
| 5.6: Maintain appropriate physical and emotional boundaries. | 614 | 2.0 | 1.9 | 1.2 | 614 | 4.0 | 3.6 | 0.7 | 0 | 1 | 0.2 |
| | | | | | | | | | 1 | 585 | 95.3 |
| | | | | | | | | | 2 | 28 | 4.6 |
| 5.7: Serve as patient advocate. | 614 | 2.0 | 2.1 | 1.3 | 614 | 4.0 | 3.6 | 0.7 | 0 | 5 | 0.8 |
| | | | | | | | | | 1 | 571 | 93.0 |
| | | | | | | | | | 2 | 38 | 6.2 |
| 5.8: Document pertinent aspects of the nurse-patient relationship in patient records. | 614 | 1.0 | 1.4 | 1.2 | 614 | 3.0 | 3.0 | 1.1 | 0 | 38 | 6.2 |
| | | | | | | | | | 1 | 538 | 87.6 |
| | | | | | | | | | 2 | 38 | 6.2 |

Domain 6 Ratings: Nurse-Patient Relationship

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|--|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| 6.1: Identify learning needs. | 613 | 2.0 | 1.8 | 1.1 | 613 | 3.0 | 3.3 | 0.8 | 0 | 3 | 0.5 |
| | | | | | | | | | 1 | 561 | 91.5 |
| | | | | | | | | | 2 | 49 | 8.0 |
| 6.2: Identify barriers to learning. | 614 | 2.0 | 1.9 | 1.1 | 614 | 3.0 | 3.3 | 0.8 | 0 | 10 | 1.6 |
| | | | | | | | | | 1 | 557 | 90.7 |
| | | | | | | | | | 2 | 47 | 7.7 |
| 6.3: Develop an individualized education plan with | 614 | 2.0 | 1.7 | 1.1 | 614 | 3.0 | 3.1 | 0.9 | 0 | 11 | 1.8 |

* Consequence Ratings: 0 = No Harm, 1 = Minimal Harm, 2 = Moderate Harm, 3 = Substantial Harm, 4 = Extreme Harm. Frequency Ratings: 0 = Never, 1 = Rarely, 2 = Sometimes, 3 = Often, 4 = Repeatedly. Performance Expectation Ratings: 0 = Not at all; 1 = Within the first 6 months of working as a <specialty> nurse; 2 = After the first 6 months of working as a <specialty> nurse.

Appendix C – Summary Statistics: Consequence, Frequency, and Performance Expectation Ratings 2010 Role Delineation Study: Gerontological Nurse – National Survey Results

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|--|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| the involvement of the patient, family, significant other, and caregiver. | | | | | | | | | 1 | 506 | 82.4 |
| | | | | | | | | | 2 | 97 | 15.8 |
| 6.4: Develop educational programs for groups. | 614 | 1.0 | 1.0 | 1.0 | 614 | 2.0 | 1.8 | 1.1 | 0 | 99 | 16.1 |
| | | | | | | | | | 1 | 214 | 34.9 |
| | | | | | | | | | 2 | 301 | 49.0 |
| 6.5: Implement the education plan. | 613 | 1.0 | 1.3 | 1.1 | 614 | 3.0 | 2.6 | 1.1 | 0 | 29 | 4.7 |
| | | | | | | | | | 1 | 422 | 68.7 |
| | | | | | | | | | 2 | 163 | 26.5 |
| 6.6: Evaluate the education plan's effectiveness. | 614 | 1.0 | 1.3 | 1.1 | 614 | 3.0 | 2.6 | 1.1 | 0 | 38 | 6.2 |
| | | | | | | | | | 1 | 402 | 65.5 |
| | | | | | | | | | 2 | 174 | 28.3 |
| 6.7: Document the education provided and its effectiveness in patient records. | 614 | 1.0 | 1.4 | 1.1 | 614 | 3.0 | 2.8 | 1.1 | 0 | 23 | 3.7 |
| | | | | | | | | | 1 | 483 | 78.7 |
| | | | | | | | | | 2 | 108 | 17.6 |

Domain 7 Ratings: Management and Leadership

| | Cons n | Cons Med | Cons Mean | Cons SD | Freq n | Freq Med | Freq Mean | Freq SD | PE Rating | PE Freq | PE Perc |
|---|-----------|-------------|--------------|------------|-----------|-------------|--------------|------------|--------------|------------|------------|
| 7.1: Serve as a preceptor. | 614 | 1.0 | 1.3 | 1.2 | 614 | 2.0 | 2.2 | 1.0 | 0 | 28 | 4.6 |
| | | | | | | | | | 1 | 169 | 27.6 |
| | | | | | | | | | 2 | 416 | 67.9 |
| 7.2: Coordinate patient safety measures. | 613 | 2.0 | 2.3 | 1.3 | 614 | 3.0 | 3.1 | 0.9 | 0 | 2 | 0.3 |
| | | | | | | | | | 1 | 477 | 77.7 |
| | | | | | | | | | 2 | 135 | 22.0 |
| 7.3: Improve quality of nursing care delivery. | 614 | 2.0 | 1.8 | 1.2 | 614 | 3.0 | 3.0 | 1.0 | 0 | 8 | 1.3 |
| | | | | | | | | | 1 | 401 | 65.3 |
| | | | | | | | | | 2 | 205 | 33.4 |
| 7.4: Serve as a clinical content expert for the design and enhancement of policies, procedures, | 613 | 1.0 | 1.4 | 1.2 | 614 | 2.0 | 2.0 | 1.1 | 0 | 65 | 10.6 |
| | | | | | | | | | 1 | 155 | 25.2 |

* Consequence Ratings: 0 = No Harm, 1 = Minimal Harm, 2 = Moderate Harm, 3 = Substantial Harm, 4 = Extreme Harm. Frequency Ratings: 0 = Never, 1 = Rarely, 2 = Sometimes, 3 = Often, 4 = Repeatedly. Performance Expectation Ratings: 0 = Not at all; 1 = Within the first 6 months of working as a <specialty> nurse; 2 = After the first 6 months of working as a <specialty> nurse.

Appendix C – Summary Statistics: Consequence, Frequency, and Performance Expectation Ratings 2010 Role Delineation Study: Gerontological Nurse – National Survey Results

| | | | | | | | | | | | | |
|--|-----|-----|-----|-----|-----|-----|-----|-----|---|-----|------|------|
| processes, and systems that affect nursing care. | | | | | | | | | | 2 | 394 | 64.2 |
| 7.5: Use electronic information systems in nursing care delivery. | 614 | 1.0 | 1.4 | 1.2 | 613 | 4.0 | 3.2 | 1.1 | 0 | 25 | 4.1 | |
| | | | | | | | | | 1 | 552 | 89.9 | |
| | | | | | | | | | 2 | 37 | 6.0 | |
| 7.6: Follow ethical standards in nursing care delivery and management. | 614 | 2.0 | 2.3 | 1.4 | 614 | 4.0 | 3.7 | 0.7 | 0 | 6 | 1.0 | |
| | | | | | | | | | 1 | 599 | 97.6 | |
| | | | | | | | | | 2 | 9 | 1.5 | |
| 7.7: Follow legal and regulatory requirements in nursing care delivery and management. | 614 | 3.0 | 2.4 | 1.4 | 614 | 4.0 | 3.7 | 0.7 | 0 | 6 | 1.0 | |
| | | | | | | | | | 1 | 590 | 96.1 | |
| | | | | | | | | | 2 | 18 | 2.9 | |

* Consequence Ratings: 0 = No Harm, 1 = Minimal Harm, 2 = Moderate Harm, 3 = Substantial Harm, 4 = Extreme Harm. Frequency Ratings: 0 = Never, 1 = Rarely, 2 = Sometimes, 3 = Often, 4 = Repeatedly. Performance Expectation Ratings: 0 = Not at all; 1 = Within the first 6 months of working as a <specialty> nurse; 2 = After the first 6 months of working as a <specialty> nurse.

**Appendix C – Summary Statistics: Consequence, Frequency, and Performance Expectation Ratings
2010 Role Delineation Study: Gerontological Nurse – National Survey Results**

Appendix D

Task Statements Sorted by Relative Importance

| Gerontological | PE Mode | Avg Cons | Avg Freq | Relative Importance |
|---|----------------|-----------------|-----------------|----------------------------|
| 1.2: Obtain a home medication list. | 1.0 | 2.46 | 3.33 | 42.5 |
| 1.5: Complete psychosocial assessment using age-appropriate, system-specific, evidence-based tools and assessment techniques. | 1.0 | 1.70 | 3.06 | 34.2 |
| 2.3: Formulate expected outcomes with patient, family, significant other, and interdisciplinary team involvement using clinical pathways or a different system. [Specify other] | 1.0 | 1.33 | 2.55 | 34.0 |
| 7.4: Serve as a clinical content expert for the design and enhancement of policies, procedures, processes, and systems that affect nursing care. | 2.0 | 1.35 | 2.00 | 33.8 |
| 3.1: Use Nursing Interventions Classification (NIC) as the basis for interventions specific to the plan of care. | 1.0 | 1.28 | 2.37 | 32.8 |
| 1.8: Review findings provided by others. | 1.0 | 1.76 | 3.10 | 32.0 |
| 6.5: Implement the education plan. | 1.0 | 1.34 | 2.56 | 32.0 |
| 7.7: Follow legal and regulatory requirements in nursing care delivery and management. | 1.0 | 2.42 | 3.73 | 30.8 |
| 4.1: Collect data related to the patient's response to interventions. | 1.0 | 1.80 | 3.29 | 29.6 |
| 1.11: Identify nursing diagnoses using clinical pathways. | 1.0 | 1.32 | 2.60 | 29.5 |
| 1.14: Document assessment findings in patient records. | 1.0 | 2.09 | 3.61 | 29.4 |
| 7.1: Serve as a preceptor. | 2.0 | 1.27 | 2.19 | 28.6 |
| 5.4: Maintain a developmentally appropriate therapeutic relationship. | 1.0 | 2.03 | 3.74 | 27.8 |
| 1.7: Obtain diagnostic test results. | 1.0 | 2.20 | 3.20 | 27.5 |
| 1.4: Perform physical examination using age-appropriate, system-specific, evidence-based assessment techniques. | 1.0 | 2.05 | 3.33 | 27.4 |
| 2.5: Document plan of care and expected outcomes in patient records. | 1.0 | 1.67 | 3.29 | 27.3 |
| 5.3: Develop rapport. | 1.0 | 1.63 | 3.69 | 26.8 |
| 3.3: Create a safe, developmentally appropriate, therapeutic environment conducive to care. | 1.0 | 2.21 | 3.48 | 26.7 |
| 3.4: Coordinate patient care. | 1.0 | 2.00 | 3.47 | 26.6 |
| 6.2: Identify barriers to learning. | 1.0 | 1.86 | 3.26 | 26.5 |
| 1.1: Obtain patient history using age-appropriate, system-specific, evidence-based tools. | 1.0 | 1.82 | 3.33 | 26.5 |
| 1.10: Identify nursing diagnoses using the North American Nursing Diagnosis Association-International (NANDA-I) taxonomy. | 1.0 | 1.21 | 2.58 | 26.0 |
| 1.13: Create a problem list based on assessment data. | 1.0 | 1.61 | 3.00 | 25.9 |
| 7.3: Improve quality of nursing care delivery. | 1.0 | 1.79 | 2.98 | 24.8 |
| 7.6: Follow ethical standards in nursing care delivery and management. | 1.0 | 2.25 | 3.67 | 24.7 |
| 2.2: Formulate expected outcomes with patient, family, significant other, and interdisciplinary team involvement using Nursing Outcomes Classification (NOC). | 1.0 | 1.36 | 2.69 | 24.4 |
| 4.3: Evaluate the patient's response to interventions and the effectiveness of the plan of care. | 1.0 | 1.93 | 3.41 | 24.3 |
| 5.7: Serve as patient advocate. | 1.0 | 2.10 | 3.60 | 24.3 |
| 4.4: Update the plan of care. | 1.0 | 1.69 | 3.25 | 24.2 |
| 6.1: Identify learning needs. | 1.0 | 1.79 | 3.29 | 24.1 |
| 6.4: Develop educational programs for groups. | 2.0 | 0.96 | 1.78 | 23.9 |
| 2.4: Develop an individualized, developmentally appropriate plan of care. | 1.0 | 1.69 | 3.14 | 22.5 |
| 3.6: Document nursing interventions in patient records. | 1.0 | 1.94 | 3.63 | 21.4 |

Appendix D – Task Statements Sorted by Relative Weight

2010 Role Delineation Study: Gerontological Nurse – National Survey Results

| Gerontological | PE Mode | Avg Cons | Avg Freq | Relative Importance |
|--|----------------|-----------------|-----------------|----------------------------|
| 6.7: Document the education provided and its effectiveness in patient records. | 1.0 | 1.38 | 2.83 | 21.3 |
| 7.5: Use electronic information systems in nursing care delivery. | 1.0 | 1.44 | 3.23 | 21.2 |
| 3.2: Use other evidence-based practice guidelines as the basis for interventions specific to the plan of care. | 1.0 | 1.53 | 2.85 | 20.9 |
| 7.2: Coordinate patient safety measures. | 1.0 | 2.31 | 3.13 | 20.4 |
| 5.2: Establish trust. | 1.0 | 1.78 | 3.72 | 20.2 |
| 1.9: Synthesize available data and knowledge to identify patterns and variances. | 1.0 | 1.65 | 2.66 | 20.0 |
| 1.12: Identify nursing diagnoses using a different system. [Specify the system] | 0.0 | 0.72 | 1.32 | 20.0 |
| 2.1: Prioritize nursing diagnoses and/or problems. | 1.0 | 1.79 | 3.30 | 19.6 |
| 6.3: Develop an individualized education plan with the involvement of the patient, family, significant other, and caregiver. | 1.0 | 1.68 | 3.08 | 19.5 |
| 4.2: Collect data on factors that impact the patient's care. | 1.0 | 1.77 | 3.13 | 19.5 |
| 6.6: Evaluate the education plan's effectiveness. | 1.0 | 1.31 | 2.57 | 19.4 |
| 5.1: Approach the patient in a developmentally appropriate manner. | 1.0 | 1.64 | 3.67 | 19.3 |
| 1.6: Collect data on psychoses. | 1.0 | 1.66 | 2.25 | 18.8 |
| 1.3: Perform a medication reconciliation | 1.0 | 2.47 | 3.26 | 18.7 |
| 4.5: Communicate changes to the patient, family, significant other, and interdisciplinary team. | 1.0 | 1.97 | 3.46 | 17.9 |
| 5.6: Maintain appropriate physical and emotional boundaries. | 1.0 | 1.89 | 3.59 | 17.3 |
| 5.8: Document pertinent aspects of the nurse-patient relationship in patient records. | 1.0 | 1.44 | 3.00 | 16.6 |
| 3.5: Administer medications as prescribed using evidence-based, developmentally appropriate, age-appropriate techniques. | 1.0 | 2.85 | 3.55 | 16.5 |
| 4.6: Document the patient's response to interventions and changes to the plan of care in patient records. | 1.0 | 1.88 | 3.46 | 16.4 |
| 5.5: Support the patient's parents, family, significant others, and caregivers. | 1.0 | 1.63 | 3.57 | 10.4 |

The American Nurses Credentialing Center (ANCC) provides individuals and organizations throughout the nursing profession with the tools they need on their journey to excellence. ANCC's internationally renowned credentialing programs certify nurses in specialty practice areas, recognize health organizations for nursing excellence through the Magnet Recognition Program®, and accredit providers and approvers of continuing nursing education. In addition, ANCC offers an array of informational and educational services and products to support its core credentialing programs.

ANCC is the largest and most prestigious nurse credentialing organization in the world. ANCC certification exams validate nurses' skills, knowledge, and abilities. More than a quarter million nurses have been certified by ANCC since 1990. More than 75,000 advanced practice nurses are currently certified by ANCC.

The ANCC Magnet Recognition Program® recognizes healthcare organizations that provide the very best in nursing care and professionalism in nursing practice. The program also provides a vehicle for disseminating best practices and strategies among nursing systems. The ANCC Magnet Recognition Program® is the gold standard for nursing excellence.

ANCC Accreditation of a continuing nursing education program is the hallmark of quality. ANCC is the nation's leader in accreditation of continuing nursing education programs.

ANCC produces a variety of educational services and products to support those seeking certification, Magnet recognition, or accreditation. Review and resource materials, review seminars, and web-based learning all support individuals who are on the path to certification. ANCC's National Magnet Conferences, workshops, and consultants support organizations on the journey to Magnet Recognition. Consultants are available to assist continuing nursing education accreditation programs in achieving the highest caliber education.

ANCC is passionate about helping nurses on their journey to nursing excellence.

American Nurses Credentialing Center (ANCC) is a subsidiary of the American Nurses Association (ANA).



8515 Georgia Avenue
Suite 400
Silver Spring, MD 20910-4392
1.800.284.2378
301.628.5000 *tel*
301.628.5004 *fax*

www.nursecredentialing.org

2010 Role Delineation Study: Gerontological Nurse – National Survey Results
© Copyright 2010 American Nurses Credentialing Center, All Rights Reserved